

TIFR/TH/97-32

June, 1997

hep-ph/9706541

χ_1 and Polarisation Asymmetries for Quarkonia at High Orders in Non-relativistic QCD

Sourendu Gupta¹ and Prakash Mathews².Theory Group, Tata Institute of Fundamental Research,
Homi Bhabha Road, Bombay 400005, India.

Abstract

We study doubly polarised asymmetries of $\bar{c}c$ and $\bar{b}b$ mesons in hadro- and photo-production at low transverse momentum in non-relativistic QCD to high orders in the relative velocity of the pair, v . We give the complete set of expressions required for the asymmetries up to order v^9 . The asymmetries in the production of $\eta_{c,b}$ states are a stable measure of the polarised gluon densities. The asymmetries for $\chi_{c,b}$, J/ψ , ψ' , and the various Υ states are stringent tests of the NRQCD scaling relations.

¹E-mail: sgupta@theory.tifr.res.in²E-mail: prakash@theory.tifr.res.in

1 Introduction

Recent progress in the understanding of cross sections for production of heavy quarkonium resonances has come through the non-relativistic QCD (NRQCD) reformulation of this problem [1]. Factorisation of the physics at a short distance scale of order $1/m$ (where m is the mass of heavy quark) from long distance scales of order $1/mv$ and $1/mv^2$ (where v is the velocity of either of the heavy quark in the rest frame of the pair) has been proven in the NRQCD formalism for processes dominated by a large transverse momentum [2]. The resulting cross sections are a double power series in v^2 and the strong coupling $\alpha_s(m^2)$. For charmonium states, a numerical coincidence, $v^2 \sim \alpha_s(m^2)$, complicates the double expansion.

The formalism has been successfully applied to large transverse momentum processes [3]. Interestingly, inclusive production cross sections for charmonium at low energies, dominated by low transverse momenta, also seem to have a good phenomenological description in terms of this approach [4, 5, 6]. The remaining phenomenological problem is to explain the χ_1/χ_2 ratio observed in hadro-production. $\chi_{0,2}$ are first produced at order $\alpha_s^2 v^5$, whereas χ_1 begins only at order $\alpha_s^2 v^9$. It was argued in [5, 6] that a better understanding of this ratio may require higher order terms in α_s and v^2 . In [7] the NRQCD series was evaluated to order $\alpha_s^2 v^9$. In the dominant gluon fusion channel a previously missed term at order $\alpha_s^2 v^7$ was found for $\chi_{0,2}$ and up to eleven terms at order $\alpha_s^2 v^9$ were obtained for all χ states.

In this paper we extend the computations of [7] to polarisation asymmetries,

$$A = \frac{\Delta\sigma}{\sigma} = \frac{\sum_{hh'} hh' \sigma(h, h')}{\sum_{hh'} \sigma(h, h')}, \quad (1.1)$$

where h and h' are the helicities of the beam and target respectively, and $\sigma(h, h')$ is the cross section for fixed initial helicities. This allows us to formulate several tests of NRQCD and the velocity scaling rules that it yields. In addition, we propose measurements of polarised gluon densities, Δg . The spin asymmetries have earlier been computed at lower orders for low transverse momentum processes [8] and at high transverse momenta [9].

The NRQCD factorisation formula for the inclusive production of heavy

quarkonium resonances H with 4-momentum P can be written as

$$\begin{aligned} d\sigma &= \frac{1}{\Phi} \frac{d^3P}{(2\pi)^3 2E_P} \sum_{ij} C_{ij} \langle \mathcal{K}_i \Pi(H) \mathcal{K}_j^\dagger \rangle, \\ d\Delta\sigma &= \frac{1}{\Phi} \frac{d^3P}{(2\pi)^3 2E_P} \sum_{ij} \tilde{C}_{ij} \langle \mathcal{K}_i \Pi(H) \mathcal{K}_j^\dagger \rangle. \end{aligned} \quad (1.2)$$

where Φ is a flux factor. The coefficient functions C_{ij} and $\tilde{C}_{ij}$ are computable in perturbative QCD and hence have an expansion in the strong coupling α_s (evaluated at the NRQCD cutoff). Although the matrix element is non-perturbative, it has a fixed scaling dimension in the quark velocity v .

The fermion bilinear operators $\mathcal{K}_i$ are built out of heavy quark fields sandwiching colour and spin matrices and the covariant derivative $\mathbf{D}$. The composite labels i and j include the colour index α , the spin quantum number S , the number of derivative operators N , the orbital angular momentum L (found by coupling the derivatives), the total angular momentum J and the helicity J_z . When the final state spin is unobserved, the hadron projection operator

$$\Pi(H) = \sum_s |H, s\rangle \langle H, s|, \quad (1.3)$$

(where s denotes hadron states with energy less than the NRQCD cutoff), is diagonal in these quantum numbers. Then it is clear that the operators $\mathcal{K}_i$ and $\mathcal{K}_j$ in eq. (1.2) are restricted to have equal L , S , J and J_z , for both $d\sigma$ and $d\Delta\sigma$.

The J_z -dependence of these matrix elements can be factored out using the Wigner-Eckart theorem—

$$\begin{aligned} \langle \mathcal{K}_i \Pi(H) \mathcal{K}_i^\dagger \rangle &= \frac{1}{2J+1} \mathcal{O}_\alpha^H(2S+1 L_J^N), \\ \langle \mathcal{K}_i \Pi(H) \mathcal{K}_j^\dagger \rangle &= \frac{1}{2J+1} \mathcal{P}_\alpha^H(2S+1 L_J^N, 2S+1 L_J^{N'}), \end{aligned} \quad (1.4)$$

where the factors of $1/(2J+1)$ come from a Clebsch-Gordan coefficient. This is conventionally included in the coefficient function. We have used the notation of [2] for both diagonal and off-diagonal operators. The NRQCD power counting rule for these matrix elements is—

$$d = 3 + N + N' + 2(E_d + 2M_d), \quad (1.5)$$

where E_d and M_d are the number of colour electric and magnetic transitions required to connect the hadronic state to the state $\mathcal{K}_i|0\rangle$.

In section 2 of this paper we present the main results for $\Delta\sigma$ after a telegraphic review of the threshold expansion technique [10], the kinematics and the appropriate Taylor series expansion of the perturbative matrix element. This material is given in greater detail in our earlier paper [7]. The phenomenology is discussed in the final section 3.

2 The Threshold Expansion

We choose to construct the coefficient functions using the “threshold expansion” technique of [10]. This consists of calculating, in perturbative QCD, the matrix element $\mathcal{M}$ connecting the initial states to final states with a heavy quark-antiquark pair ($\bar{Q}Q$), and Taylor expanding the result in the relative momentum of the pair, q , after performing a non-relativistic reduction of the Dirac spinors. The resulting expression is squared and matched to the NRQCD formula of eq. (1.2) by inserting a perturbative projector onto a non-relativistic $\bar{Q}Q$ state between the two spinor bilinears. The coefficient of this matrix element is the required coefficient function.

Symbolically—

$$\begin{aligned}\frac{1}{4}\sum_{hh'}|\mathcal{M}(h,h')|^2 &= \sum_{ij}C_{ij}\langle\mathcal{K}_i\Pi(\bar{Q}Q)\mathcal{K}_j^\dagger\rangle, \\ \frac{1}{4}\sum_{hh'}hh'|\mathcal{M}(h,h')|^2 &= \sum_{ij}\tilde{C}_{ij}\langle\mathcal{K}_i\Pi(\bar{Q}Q)\mathcal{K}_j^\dagger\rangle,\end{aligned}\tag{2.1}$$

where the left hand sides are Taylor expanded in q . Each factor of q Fourier transforms into a factor of the covariant derivative $\mathbf{D}$ on the right hand side. Since each matrix element on the right of eq. (2.1) corresponds to a unique matrix element in eq. (1.2), the order up to which the Taylor expansion is to be performed is determined by the scaling of the non-perturbative matrix elements with v . Since we require a classification of operators by the angular momentum, it turns out to be very convenient to use the spherical tensor methods detailed in [7].

In this paper we evaluate the polarised cross sections to order $\alpha_s^2 v^9$. The Taylor expansion order, $N+N'\leq 6$ is obtained by setting $d=9$ and $E_d=M_d=0$ in eq. (1.5). Furthermore, since we examine the leading term in




Figure 1: The ratios $\mathcal{L}_{\bar{q}q}/\mathcal{L}_{gg}$ (full line) and $\Delta\mathcal{L}_{\bar{q}q}/\mathcal{L}_{gg}$ (broken line) computed using the GRSV LO parton density set [11] as a function of the CM energy $\sqrt{S}$ at zero rapidity.

perturbation theory, the perturbative projector has only one term—

$$\Pi(\bar{Q}Q) = |\bar{Q}Q\rangle\langle\bar{Q}Q|. \quad (2.2)$$

In agreement with [7, 10] we use the relativistic normalisation of states

$$\langle Q(p, \xi) \bar{Q}(q, \eta) | Q(p', \xi') \bar{Q}(q', \eta') \rangle = 4E_p E_q (2\pi)^6 \delta^3(p - p') \delta^3(q - q'), \quad (2.3)$$

with the spinor normalisations $\xi^\dagger \xi = \eta^\dagger \eta = 1$. Expanding $E_p = E_q = \sqrt{m^2 + q^2}$ in q^2 allows us to write the spinor bilinears in terms of transition operators built out of the heavy quark field. The kinematics is very simple to leading order in α_s . The momenta of the initial particles are p_1 and p_2 . We take p_1 to lie in the positive z -direction and p_2 to be oppositely directed. The net momentum $P = p_1 + p_2$.

The 4-momenta of Q and $\bar{Q}$ (p and $\bar{p}$ respectively) are written as

$$p = \frac{1}{2}P + L_j q^j \quad \text{and} \quad \bar{p} = \frac{1}{2}P - L_j q^j. \quad (2.4)$$

Note that $p^2 = \bar{p}^2 = m^2$, where m is the mass of the heavy quark. The space-like vector q is always defined in the rest frame of the pair, and L_j^μ

boosts it to any frame. We shall use Greek indices for Lorentz tensors and Latin indices for Euclidean 3-tensors.

In the next two sections we will write down the parton level cross sections, $\Delta\hat{\sigma}$ for $\bar{q}q \rightarrow H$ and $gg \rightarrow H$ (where H is a quarkonium state). The hadronic cross sections, $\Delta\sigma$, are obtained by multiplying these by appropriate parton luminosities—

$$\Delta\mathcal{L}_{\bar{q}q} = \sum_f \Delta q_f(x_1) \Delta \bar{q}_f(x_2), \quad \Delta\mathcal{L}_{gg} = \Delta g(x_1) \Delta g(x_2), \quad (2.5)$$

where $x_1 = (2m/\sqrt{S}) \exp(y)$ and $x_2 = (2m/\sqrt{S}) \exp(-y)$, $\sqrt{S}$ is the centre of mass (CM) energy of the colliding protons, y is the rapidity at which the quarkonium is observed, and $\Delta q_f(x)$, $\Delta \bar{q}_f(x)$ and $\Delta g(x)$ are the quark, antiquark (of flavour f) and gluon polarised densities. For unpolarised cross sections, σ , we must similarly define unpolarised luminosities $\mathcal{L}_{\bar{q}q}$ and $\mathcal{L}_{gg}$ using the corresponding unpolarised parton densities. As shown in Figure 1, for $\sqrt{S} \geq 20$ GeV, $\mathcal{L}_{\bar{q}q} \ll \mathcal{L}_{gg}$. Also, present day data indicates that $|\Delta\mathcal{L}_{\bar{q}q}| \ll \mathcal{L}_{gg}$. Consequently, the $\bar{q}q$ channel may be neglected for double polarised asymmetries to good precision.

2.1 $\bar{q}q \rightarrow \bar{Q}Q$

We write down the polarised matrix element for the subprocess $\bar{q}q \rightarrow \bar{Q}Q$ for completeness, and because it provides a simple illustration of the techniques used. It is given exactly by the expression

$$\begin{aligned} \mathcal{M} = & -\frac{ig^2}{M^2} [\bar{v}(p_2, h') \gamma_\mu T^a u(p_1, h)] L_j^\mu \\ & \times \left[M \xi^\dagger \sigma^j T^a \eta - \frac{4}{M + 2m} q^j \xi^\dagger (q \cdot \sigma) T^a \eta \right], \end{aligned} \quad (2.6)$$

where T^a is a colour generator, u and v are the light quark spinors with helicities h and h' respectively, and ξ and η are the heavy quark Pauli spinors. The equations of motion for the initial state quarks have been used to obtain the explicitly gauge invariant matrix element above. The desired Taylor series expansion is obtained by using the relation $M^2 = 4(m^2 + q^2)$ to expand all factors with M .

The squared matrix element for this process is easily written down. The difference between the polarised and unpolarised cross sections follows when

one notices that the light quark helicities give an overall factor of $1 - hh'$ in the squared matrix element. As a result, $\tilde{C}_{ij} = -C_{ij}$ and the parton level spin asymmetries are all -1 . We list the parton level polarised cross sections—

$$\begin{aligned}
\Delta\hat{\sigma}_{\bar{q}q}^{\eta_c} &= -\frac{\pi^3\alpha_s^2}{54m^4}\delta(\hat{s}-4m^2)\mathcal{O}_8^{\eta_c}({}^3S_1^0), \\
\Delta\hat{\sigma}_{\bar{q}q}^{h_c} &= -\frac{\pi^3\alpha_s^2}{54m^4}\delta(\hat{s}-4m^2)\mathcal{O}_8^{h_c}({}^3S_1^0), \\
\Delta\hat{\sigma}_{\bar{q}q}^{J/\psi} &= -\frac{\pi^3\alpha_s^2}{54m^4}\delta(\hat{s}-4m^2)\left[\mathcal{O}_8^{J/\psi}({}^3S_1^0)\right. \\
&\quad \left.+\frac{1}{m^2}\left\{\frac{2}{\sqrt{3}}\mathcal{P}_8^{J/\psi}({}^3S_1^0, {}^3S_1^2)+\frac{1}{4}\mathcal{O}_8^{J/\psi}({}^3P_1^2)\right\}\right], \\
\Delta\hat{\sigma}_{\bar{q}q}^{\chi_J} &= -\frac{\pi^3\alpha_s^2}{54m^4}\delta(\hat{s}-4m^2)\left[\mathcal{O}_8^{\chi_J}({}^3S_1^0)+\frac{2}{\sqrt{3}m^2}\mathcal{P}_8^{\chi_J}({}^3S_1^0, {}^3S_1^2)\right. \\
&\quad \left.+\frac{1}{m^4}\left\{\frac{4}{3}\mathcal{O}_8^{\chi_J}({}^3S_1^2)+\frac{5}{12}\mathcal{O}_8^{\chi_J}({}^3D_1^2)+\frac{7\sqrt{5}}{12}\mathcal{P}_8^{\chi_J}({}^3S_1^0, {}^3S_1^4)\right\}\right],
\end{aligned} \tag{2.7}$$

where $\hat{s}$ is the parton CM energy.

2.2 $gg \rightarrow \bar{Q}Q$

The squared matrix element for the gg process is technically a little more complicated. We work in a class of ghost-free gauges called planar gauges [12]. The density matrix for an initial state gluon of momentum p and helicity h in these gauges is given by

$$\epsilon_\mu(p, h)\epsilon_\nu^*(p, h) = \frac{1}{2}\left[-g_{\mu\nu} + \frac{1}{p \cdot V}(p_\mu V_\nu + p_\nu V_\mu) + \frac{ih}{p \cdot V}\epsilon_{\mu\nu\rho\sigma}p^\rho V^\sigma\right]. \tag{2.8}$$

The vector V defines the gauge choice. We write $V = c_1 p_1 + c_2 p_2$, with $c_1/c_2 \sim \mathcal{O}(1)$. We verify that all our results are gauge invariant by the explicit check that they do not depend on the arbitrary coefficients c_1 and c_2 .

The sum of the matrix elements arising from the three Feynman diagrams (s -channel gluon exchange, $\mathcal{M}_s$, and t - and u -channel quark exchanges, $\mathcal{M}_t$ and $\mathcal{M}_u$) can be decomposed into three colour amplitudes—

$$\mathcal{M} = \frac{1}{6}g^2\delta_{ab}S + \frac{1}{2}g^2d_{abc}D^c + \frac{i}{2}g^2f_{abc}F^c. \tag{2.9}$$

The colour amplitudes S and D involve only $\mathcal{M}_t + \mathcal{M}_u$, whereas F involves $\mathcal{M}_s$ as well as $\mathcal{M}_t - \mathcal{M}_u$.

In order to write down our results, we find it convenient to introduce the notation

$$\mathcal{A} = \frac{1}{M^2} \varepsilon_{\lambda\sigma\mu\nu} p_1^\lambda p_2^\sigma \epsilon_1^\mu \epsilon_2^\nu \quad \text{and} \quad \mathcal{S}_{ij} = A_i \hat{z}_j + A_j \hat{z}_i - B_{ij} + \epsilon_1 \cdot \epsilon_2 \hat{z}_i \hat{z}_j, \quad (2.10)$$

where

$$\begin{aligned} A_i &= \frac{1}{M} (\epsilon_1 \cdot L_i \epsilon_2 \cdot p_1 - \epsilon_2 \cdot L_i \epsilon_1 \cdot p_2), \\ B_{ij} &= \epsilon_1 \cdot L_i \epsilon_2 \cdot L_j + \epsilon_2 \cdot L_i \epsilon_1 \cdot L_j. \end{aligned} \quad (2.11)$$

Here ϵ_i is the polarisation vector for the initial gluon of momentum p_i . In order to identify all terms to order v^9 we need the colour amplitude S to order q^5 —

$$\begin{aligned} S &= -\left(\frac{8im}{M}\right) \mathcal{A}(\xi^\dagger \eta) + \frac{4}{M} \mathcal{S}_{jm} (q^m \xi^\dagger \sigma^j \eta) - \left(\frac{32im}{M^3}\right) \mathcal{A} \hat{z}_m \hat{z}_n (q^m q^n \xi^\dagger \eta) \\ &\quad + \frac{16}{M^3} \left[\mathcal{S}_{jm} \hat{z}_n \hat{z}_p - \frac{M}{M+2m} \delta_{jm} \mathcal{S}_{np} \right] (q^m q^n q^p \xi^\dagger \sigma^j \eta) \\ &\quad - \left(\frac{128im}{M^5}\right) \mathcal{A} \hat{z}_m \hat{z}_n \hat{z}_p \hat{z}_r (q^m q^n q^p q^r \xi^\dagger \eta) \\ &\quad + \frac{64}{M^5} \left[\mathcal{S}_{jm} \hat{z}_n \hat{z}_p - \frac{M}{M+2m} \delta_{jm} \mathcal{S}_{np} \right] \hat{z}_r \hat{z}_s (q^m q^n q^p q^r q^s \xi^\dagger \sigma^j \eta), \end{aligned} \quad (2.12)$$

where $\hat{z}$ is the unit vector in the z -direction. The amplitude D differs only through having colour octet matrix elements in place of the colour singlet ones shown above. For the colour amplitude F we need the expansion

$$F^c = -\left(\frac{16im}{M^2}\right) \mathcal{A} \hat{z}_m (q^m \xi^\dagger T^c \eta) + \frac{8}{M^2} \mathcal{S}_{jm} \hat{z}_n (q^m q^n \xi^\dagger \sigma^j T^c \eta). \quad (2.13)$$

In all three colour amplitudes, the terms in $\mathcal{A}$ are spin singlet and those in $\mathcal{S}$ are spin triplet.

The density matrix in eq. (2.8) yields

$$\begin{aligned} \Delta \mathcal{S} \cdot \mathcal{S}^* &\equiv \frac{1}{4} \sum_{hh'} hh' S_{jm} S_{j'm'}^* = \frac{3}{2} [\sigma q]_0^0 [\sigma^\dagger q^\dagger]_0^0 - \sum_{\lambda=\pm 2} [\sigma q]_\lambda^2 [\sigma^\dagger q^\dagger]_{-\lambda}^2, \\ \mathcal{S} \cdot \mathcal{S}^* &\equiv \frac{1}{4} \sum_{hh'} S_{jm} S_{j'm'}^* = \frac{3}{2} [\sigma q]_0^0 [\sigma^\dagger q^\dagger]_0^0 + \sum_{\lambda=\pm 2} [\sigma q]_\lambda^2 [\sigma^\dagger q^\dagger]_{-\lambda}^2. \end{aligned} \quad (2.14)$$

We also find, $\mathcal{A} \cdot \mathcal{A}^* = \Delta \mathcal{A} \cdot \mathcal{A}^* = 1/8$ and $\mathcal{A} \cdot \mathcal{S}^* = \Delta \mathcal{A} \cdot \mathcal{S}^* = 0$. Consequently, the difference between the unpolarised [7] and polarised cases lies solely in the flipped sign of the $J = 2$ part in eq. (2.14). Furthermore, even and odd terms in the three colour amplitudes of eq. (2.9) do not interfere with each other.

The direct 3S_1 polarised subprocess cross section is

$$\Delta \hat{\sigma}_{gg}^{J/\psi}(\hat{s}) = \varphi \left[\frac{5}{48} \tilde{\Theta}_D^{J/\psi}(7) + \left\{ \frac{5}{48} \tilde{\Theta}_D^{J/\psi}(9) + \frac{3}{16} \tilde{\Theta}_F^{J/\psi}(9) \right\} \right], \quad (2.15)$$

where

$$\varphi = \frac{\pi^3 \alpha_s^2}{4m^2} \delta(\hat{s} - 4m^2), \quad (2.16)$$

and $\tilde{\Theta}_a^{J/\psi}(d)$ denotes combinations of non-perturbative matrix elements from the colour amplitude a ($= S, D$ or F) at order v^d for the polarised subprocess cross section. These are given by

$$\begin{aligned} \tilde{\Theta}_D^{J/\psi}(7) &= \frac{1}{2m^2} \mathcal{O}_8^{J/\psi}(^1S_0^0) + \frac{1}{2m^4} \left[3\mathcal{O}_8^{J/\psi}(^3P_0^1) - \frac{4}{5} \mathcal{O}_8^{J/\psi}(^3P_2^1) \right], \\ \tilde{\Theta}_D^{J/\psi}(9) &= \frac{1}{\sqrt{3}m^4} \mathcal{P}_8^{J/\psi}(^1S_0^0, ^1S_0^2) + \frac{1}{\sqrt{15}m^6} \left[\frac{35}{4} \mathcal{P}_8^{J/\psi}(^3P_0^1, ^3P_0^3) \right. \\ &\quad \left. - 2\mathcal{P}_8^{J/\psi}(^3P_2^1, ^3P_2^3) \right], \\ \tilde{\Theta}_F^{J/\psi}(9) &= \frac{1}{2m^6} \left[\frac{1}{3} \mathcal{O}_8^{J/\psi}(^3P_1^2) + \frac{2}{5} \mathcal{O}_8^{J/\psi}(^3P_2^2) \right]. \end{aligned} \quad (2.17)$$

As a consequence of eq. (2.14) the coefficient of $J = 2$ matrix elements changes sign between the polarised and unpolarised cases. The cross sections for ψ' , Υ and all other 3S_1 states need only the replacement of the appropriate matrix elements in eq. (2.17).

The 3P_0 polarised subprocess cross section is

$$\begin{aligned} \Delta \hat{\sigma}_{gg}^{\chi_0}(\hat{s}) &= \varphi \left[\frac{1}{18} \tilde{\Theta}_S^{\chi_0}(5) + \frac{1}{18} \tilde{\Theta}_S^{\chi_0}(7) \right. \\ &\quad \left. + \left\{ \frac{1}{18} \tilde{\Theta}_S^{\chi_0}(9) + \frac{5}{48} \tilde{\Theta}_D^{\chi_0}(9) + \frac{3}{16} \tilde{\Theta}_F^{\chi_0}(9) \right\} \right]. \end{aligned} \quad (2.18)$$

where combinations of non-perturbative matrix elements appearing in eq. (2.18) are

$$\begin{aligned}
\tilde{\Theta}_S^{\chi_0}(5) &= \frac{3}{2m^4} \mathcal{O}_1^{\chi_0}(^3P_0^1), \\
\tilde{\Theta}_S^{\chi_0}(7) &= \frac{7\sqrt{5}}{4\sqrt{3}m^6} \mathcal{P}_1^{\chi_0}(^3P_0^1, ^3P_0^3), \\
\tilde{\Theta}_S^{\chi_0}(9) &= \frac{1}{8m^8} \left[\frac{245}{9} \mathcal{O}_1^{\chi_0}(^3P_0^3) + \frac{149\sqrt{7}}{10\sqrt{3}} \mathcal{P}_1^{\chi_0}(^3P_0^1, ^3P_0^5) \right] - \frac{2}{5m^4} \mathcal{O}_1^{\chi_0}(^3P_2^1), \\
\tilde{\Theta}_D^{\chi_0}(9) &= \frac{1}{2m^2} \mathcal{O}_8^{\chi_0}(^1S_0^0) + \frac{1}{2m^4} \left[3\mathcal{O}_8^{\chi_0}(^3P_0^1) - \frac{4}{5} \mathcal{O}_8^{\chi_0}(^3P_2^1) \right], \\
\tilde{\Theta}_F^{\chi_0}(9) &= \frac{1}{6m^4} \mathcal{O}_8^{\chi_0}(^1P_1^1) + \frac{1}{18m^6} \left[\mathcal{O}_8^{\chi_0}(^3S_1^2) + 5\mathcal{O}_8^{\chi_0}(^3D_1^2) \right].
\end{aligned} \tag{2.19}$$

The difference between these and the unpolarised cross sections of [7] are only in the sign of the coefficient of the $J = 2$ components in the combinations $\tilde{\Theta}_S^{\chi_0}(9)$ and $\tilde{\Theta}_D^{\chi_0}(9)$.

For the 3P_1 state we find

$$\Delta\hat{\sigma}_{gg}^{\chi_1}(\hat{s}) = \varphi \left[\frac{1}{18} \tilde{\Theta}_S^{\chi_1}(9) + \frac{5}{48} \tilde{\Theta}_D^{\chi_1}(9) + \frac{3}{16} \tilde{\Theta}_F^{\chi_1}(9) \right], \tag{2.20}$$

where the combinations of non-perturbative matrix elements are,

$$\begin{aligned}
\tilde{\Theta}_S^{\chi_1}(9) &= \frac{1}{2m^4} \left[3\mathcal{O}_1^{\chi_1}(^3P_0^1) - \frac{4}{5} \mathcal{O}_1^{\chi_1}(^3P_2^1) \right], \\
\tilde{\Theta}_D^{\chi_1}(9) &= \frac{1}{2m^2} \mathcal{O}_8^{\chi_1}(^1S_0^0) + \frac{1}{2m^4} \left[3\mathcal{O}_8^{\chi_1}(^3P_0^1) - \frac{4}{5} \mathcal{O}_8^{\chi_1}(^3P_2^1) \right], \\
\tilde{\Theta}_F^{\chi_1}(9) &= \frac{1}{6m^4} \mathcal{O}_8^{\chi_1}(^1P_1^1) + \frac{1}{3m^6} \left[\frac{1}{6} \mathcal{O}_8^{\chi_1}(^3S_1^2) \right. \\
&\quad \left. + \frac{5}{6} \mathcal{O}_8^{\chi_1}(^3D_1^2) + \frac{1}{5} \mathcal{O}_8^{\chi_1}(^3D_2^2) \right].
\end{aligned} \tag{2.21}$$

The 3P_2 cross section is

$$\begin{aligned}
\Delta\hat{\sigma}_{gg}^{\chi_2}(\hat{s}) &= \varphi \left[\frac{1}{18} \tilde{\Theta}_S^{\chi_2}(5) + \frac{1}{18} \tilde{\Theta}_S^{\chi_2}(7) \right. \\
&\quad \left. + \left\{ \frac{1}{18} \tilde{\Theta}_S^{\chi_2}(9) + \frac{5}{48} \tilde{\Theta}_D^{\chi_2}(9) + \frac{3}{16} \tilde{\Theta}_F^{\chi_2}(9) \right\} \right]
\end{aligned} \tag{2.22}$$

where the combinations of non-perturbative matrix elements are

$$\begin{aligned}
\tilde{\Theta}_S^{\chi_2}(5) &= -\frac{2}{5m^4}\mathcal{O}_1^{\chi_2}({}^3P_2^1), \\
\tilde{\Theta}_S^{\chi_2}(7) &= -\frac{2}{\sqrt{15}m^6}\mathcal{P}_1^{\chi_2}({}^3P_2^1, {}^3P_2^3), \\
\tilde{\Theta}_S^{\chi_2}(9) &= \frac{3}{2m^4}\mathcal{O}_1^{\chi_2}({}^3P_0^1) - \frac{1}{75m^8}\left[\frac{238}{9}\mathcal{O}_1^{\chi_2}({}^3P_2^3) + \frac{141\sqrt{3}}{2\sqrt{7}}\mathcal{P}_1^{\chi_2}({}^3P_2^1, {}^3P_2^5)\right], \\
\tilde{\Theta}_D^{\chi_2}(9) &= \frac{1}{2m^2}\mathcal{O}_8^{\chi_2}({}^1S_0^0) + \frac{1}{2m^4}\left[3\mathcal{O}_8^{\chi_2}({}^3P_0^1) - \frac{4}{5}\mathcal{O}_8^{\chi_2}({}^3P_2^1)\right], \\
\tilde{\Theta}_F^{\chi_2}(9) &= \frac{1}{6m^4}\mathcal{O}_8^{\chi_2}({}^1P_1^1) + \frac{1}{3m^6}\left[\frac{1}{6}\mathcal{O}_8^{\chi_2}({}^3S_1^2) + \frac{5}{6}\mathcal{O}_8^{\chi_2}({}^3D_1^2) \right. \\
&\quad \left. + \frac{1}{5}\mathcal{O}_8^{\chi_2}({}^3D_2^2) - \frac{2}{7}\mathcal{O}_8^{\chi_2}({}^3D_3^2)\right].
\end{aligned} \tag{2.23}$$

In the combination $\tilde{\Theta}_S^{\chi_2}(9)$, the coefficient of $\mathcal{O}_1^{\chi_2}({}^3P_2^3)$ is different from the unpolarised in magnitude as well as sign, because it arises from several sources. Only the part from the combination $\Delta\mathcal{S} \cdot \mathcal{S}^*$ flips sign. The other $J = 2$ terms change sign as expected. The $J = 3$ term in the F colour amplitude is also derived from the $J = 2$ part of the $\Delta\mathcal{S} \cdot \mathcal{S}^*$ term, and hence flips sign.

The production cross section for 1S_0 states is—

$$\begin{aligned}
\Delta\hat{\sigma}_{gg}^\eta(\hat{s}) &= \varphi\left[\frac{1}{18}\tilde{\Theta}_S^\eta(3) + \frac{1}{18}\tilde{\Theta}_S^\eta(5) \right. \\
&\quad \left. + \left\{\frac{1}{18}\tilde{\Theta}_S^\eta(7) + \frac{5}{48}\tilde{\Theta}_D^\eta(7) + \frac{3}{16}\tilde{\Theta}_F^\eta(7)\right\}\right],
\end{aligned} \tag{2.24}$$

where the combinations of non-perturbative matrix elements are

$$\begin{aligned}
\tilde{\Theta}_S^\eta(3) &= \frac{1}{2m^2}\mathcal{O}_1^\eta({}^1S_0^0), \\
\tilde{\Theta}_S^\eta(5) &= \frac{1}{\sqrt{3}m^4}\mathcal{P}_1^\eta({}^1S_0^0, {}^1S_0^2), \\
\tilde{\Theta}_S^\eta(7) &= \frac{1}{3m^6}\left[2\mathcal{O}_1^\eta({}^1S_0^2) + \frac{4}{\sqrt{5}}\mathcal{P}_1^\eta({}^1S_0^0, {}^1S_0^4)\right], \\
\tilde{\Theta}_D^\eta(7) &= \frac{1}{2m^2}\mathcal{O}_8^\eta({}^1S_0^0), \\
\tilde{\Theta}_F^\eta(7) &= \frac{1}{6m^4}\mathcal{O}_8^{\eta_c}({}^1P_1^1).
\end{aligned} \tag{2.25}$$

There is no $J = 2$ contribution to η_c to $\mathcal{O}(v^7)$ and hence the parton level asymmetry is identically equal to 1.

The cross section for the 1P_1 quarkonium state is—

$$\Delta\hat{\sigma}_{gg}^h(\hat{s}) = \varphi \left[\frac{5}{48}\tilde{\Theta}_D^h(5) + \frac{5}{48}\tilde{\Theta}_D^h(7) + \left\{ \frac{1}{18}\tilde{\Theta}_S^h(9) + \frac{5}{48}\tilde{\Theta}_D^h(9) + \frac{3}{16}\tilde{\Theta}_F^h(9) \right\} \right], \quad (2.26)$$

where the combinations of non-perturbative matrix elements is given by

$$\begin{aligned} \tilde{\Theta}_D^h(5) &= \frac{1}{2m^2}\mathcal{O}_8^h(^1S_0^0), \\ \tilde{\Theta}_D^h(7) &= \frac{1}{\sqrt{3}m^4}\mathcal{P}_8^h(^1S_0^0, ^1S_0^2), \\ \tilde{\Theta}_S^h(9) &= \frac{1}{2m^2}\mathcal{O}_1^h(^1S_0^0), \\ \tilde{\Theta}_D^h(9) &= \frac{1}{3m^6} \left[2\mathcal{O}_8^h(^1S_0^2) + \frac{4}{\sqrt{5}}\mathcal{P}_8^h(^1S_0^0, ^1S_0^4) \right] \\ &\quad + \frac{1}{2m^4} \left[3\mathcal{O}_8^h(^3P_0^1) - \frac{4}{5}\mathcal{O}_8^h(^3P_2^1) \right] + \frac{1}{15m^6}\mathcal{O}_8^h(^1D_2^2), \\ \tilde{\Theta}_F^h(9) &= \frac{1}{6m^4}\mathcal{O}_8^h(^1P_1^1). \end{aligned} \quad (2.27)$$

The coefficient of the matrix element $\mathcal{O}_8^h(^3P_2^1)$ in $\tilde{\Theta}_D^h(9)$ flips sign as expected. However, the other $J = 2$ term, $\mathcal{O}_8^h(^1D_2^2)$, is unaffected as it comes from $\Delta\mathcal{A} \cdot \mathcal{A}^*$.

2.3 $\gamma g \rightarrow \bar{Q}Q$ and $\gamma\gamma \rightarrow \bar{Q}Q$

The matrix elements for the two processes $\gamma p \rightarrow \bar{Q}Q$ and $\gamma\gamma \rightarrow \bar{Q}Q$ are closely related to the gg amplitudes. It is easy to check that

$$\mathcal{M}_{\gamma g} = geD, \quad \text{and} \quad \mathcal{M}_{\gamma\gamma} = e^2S, \quad (2.28)$$

where D and S are the colour amplitudes given in eq. (2.9), and e is the charge of the heavy quark.

The γg cross sections for the production of any quarkonium state can be obtained from those for the gg process, eqs. (2.15)–(2.27), by the following

prescription— replace α_s^2 in φ (eq. 2.16) by $\alpha\alpha_s$, delete the $\tilde{\Theta}_S$ and $\tilde{\Theta}_F$ terms, and replace the colour factor $5/48$ for the terms in $\tilde{\Theta}_D$ by 2 .

These computations may be applied to double polarised asymmetries in almost elastic photo-production of quarkonium states. Then the hadron level cross sections are obtained from these parton level quantities by multiplying the latter by the unpolarised or polarised gluon densities, $g(x)$ and $\Delta g(x)$ respectively. Here $x = 2m^2/m_N E_\gamma$, m_N is the proton mass and E_γ is the photon energy in the rest frame of the proton. In phenomenological applications the diffractive parts of these cross sections have to be separated. Techniques have been developed at the HERA experiments to do this [17].

The $\gamma\gamma$ cross sections are obtained with the prescription— replace α_s^2 in φ (eq. 2.16) by α^2 , delete the $\tilde{\Theta}_D$ and $\tilde{\Theta}_F$ terms in eqs. (2.15)–(2.27), and replace the colour factor $1/18$ for the terms in $\tilde{\Theta}_S$ by 16 . No parton densities enter for point-like photon cross sections. If, on the other hand, the cross sections are dominated by resolved photon processes, then the calculation is the same as for hadro-production.

3 Phenomenology

In spite of the large number of non-perturbative matrix elements in the final results, it is possible to make several quantitative and qualitative comments about the polarisation asymmetries. The most reliable predictions are obtained when the parton level asymmetries are ± 1 (the 1S_0 and 1P_1 quarkonium states) and the matrix elements drop out of the asymmetries. In these cases the polarisation asymmetries may be used to measure the polarised gluon densities. In several other cases large simplifications occur. Some of the asymmetries can be used to test NRQCD or the scaling laws for matrix elements.

Many of the matrix elements involved in the phenomenology are unknown. We develop a scaling argument which allows us to make rough estimates. A dimensional argument, neglecting possible logarithms of m and v , can be used to write

$$\langle \mathcal{K}_i \Pi(H) \mathcal{K}_j^\dagger \rangle = R_H Y_{ij} \Lambda^{D_{ij}} v^d, \quad (3.1)$$

where D_{ij} is the mass dimension of the operator, d is the velocity scaling exponent in NRQCD (eq. 1.5), Λ is the cutoff scale below which NRQCD is defined ($\Lambda \sim m$), and Y_{ij} and R_H are dimensionless numbers. We have

Matrix Elements	Values	R_H
$\mathcal{O}_8^{J/\psi} (^3S_1)$	$(1.12 \pm 0.14) \times 10^{-2} \text{ GeV}^3$	0.22 ± 0.03
$\mathcal{O}_8^{J/\psi} (^1S_0) + \frac{7}{2m^2} \mathcal{O}_8^{J/\psi} (^3P_0)$	$(3.90 \pm 1.14) \times 10^{-2} \text{ GeV}^3$	0.17 ± 0.05
$\mathcal{O}_8^{\psi'} (^3S_1)$	$(0.46 \pm 0.08) \times 10^{-2} \text{ GeV}^3$	0.09 ± 0.02
$\mathcal{O}_8^{\psi'} (^1S_0) + \frac{7}{2m^2} \mathcal{O}_8^{\psi'} (^3P_0)$	$(1.60 \pm 0.51) \times 10^{-2} \text{ GeV}^3$	0.07 ± 0.02

Table 1: The values of various non-perturbative matrix elements in the non-relativistic normalisation [13]. These can be converted to the relativistic normalisation by multiplying by $4m$. The dimensionless number R_H (eq. 3.1) has been extracted using $\Lambda = m = 1.5 \text{ GeV}$, $v^2 = 0.3$, and $Y = 1$, except in the 2nd and 4th lines of the table where we used $Y = 4.5$.

separated these two factors in order to specify that Y_{ij} contains all trivial numerical factors as well as Clebsch-Gordan coefficients, for example those which come from the approximate heavy-quark spin symmetry. R_H contains the irreducible minimum non-perturbative information. In the context of potential models, R_H would be related to some integral of the radial wave function of the state H . The scaling rules of NRQCD are a good guide to the importance of a given term if R_H depends mainly on H and is approximately independent of the operators $\mathcal{K}_{i,j}$.

In Table 1 we collect the values of some of the non-perturbative matrix elements quoted in the literature and extract from them the values of R_H . In [13] the colour octet matrix elements were fitted to collider data. We have quoted the values obtained using the GRV LO parton densities. The good agreement between the values of R_H extracted from the two sets of octet matrix elements, for both $H = J/\psi$ and ψ' , provide some basis for the working hypothesis that R_H is almost independent of the operator. In this section we shall accept it, keeping in mind that further checks are necessary.

Assuming the constancy of R_H and using heavy-quark symmetry, we find that the non-perturbative matrix elements contribute approximately $12R_\chi m^2 v^9$ to the χ_1 cross section and about $(1 + v^2 + 12v^4)R_\chi m^2 v^5$ to the

χ_2 cross sections. Then we expect

$$\frac{\sigma(\chi_1)}{\sigma(\chi_2)} \approx \frac{12v^4}{1 + v^2 + 12v^4} = 0.45, \quad (3.2)$$

independent of $\sqrt{S}$. This estimate³ is in reasonable agreement with the measured values in proton-nucleon collisions— 0.34 ± 0.16 at $\sqrt{S} = 38.8$ GeV [14] and 0.24 ± 0.28 at $\sqrt{S} = 19.4$ [15]. The measurements are also compatible with a lack of $\sqrt{S}$ dependence. In NRQCD this ratio cannot depend on the beam hadron. It turns out that the estimate in eq. (3.2) is not very far from the recently measured value in pion-nucleon collisions— 0.57 ± 0.19 at $\sqrt{S} = 31.1$ GeV [16]. However, the experimental situation certainly needs clarification.

This paper is concerned with low transverse momentum production. Our results are best suited to help design fixed target polarised experiments, for example by extracting the RHIC or HERA polarised beams for use on polarised targets. The reason is that the polarisation asymmetries are likely to be largest at small $\sqrt{S}$ (Figure 2), yielding the best signals. Also, at smaller values of $\sqrt{S}$ the $\bar{q}q$ channel seems to be unimportant for the phenomenology, and the polarised gluon asymmetries can be studied directly. Since many aspects of low-energy quarkonium production can be tested by such experiments, it is worth considering them seriously.

For 1S_0 quarkonium production, the parton level asymmetry in the gg channel is identically 1. For the yet to be discovered 1P_1 resonances, this is true with corrections of order v^4 (from the $\tilde{\Theta}_D^h(9)$ term in eq. 2.27). Only the colour amplitude S enters in $\gamma\gamma$ reactions; hence the polarisation asymmetry is exactly 1 for both these states. In γp collisions

$$A_{\gamma p} = \frac{\Delta g(x)}{g(x)} \quad (^1S_0, ^1P_1 \text{ quarkonia}), \quad (3.3)$$

For 1S_0 states the above equation is exact, whereas there are order v^4 corrections for 1P_1 states.

³The number of terms allowed at higher orders in v can grow at most by 12 for each order. If the coefficient functions continue to be of order unity, then the all orders result for the χ_1/χ_2 ratio can be bounded by 0.67 for χ_c and by 0.12 for χ_b . The ratio in photo-production is close to unity in both these cases. In [5] an estimate of $\mathcal{O}(\alpha_s^3)$ effects was used to show that the χ_1/χ_2 ratio could be about 0.3.




Figure 2: The asymmetries (a) $\Delta\mathcal{L}_{gg}/\mathcal{L}_{gg}$ (full line) (b) $(\Delta\mathcal{L}_{gg} + \Delta\mathcal{L}_{\bar{q}q})/(\mathcal{L}_{gg} + \mathcal{L}_{\bar{q}q})$ (dashed line) and (c) $(\Delta\mathcal{L}_{gg} - \Delta\mathcal{L}_{\bar{q}q})/(\mathcal{L}_{gg} + \mathcal{L}_{\bar{q}q})$ (dotted line) for zero rapidity as a function of the CM energy $\sqrt{S}$ using the GRSV LO parton density set [11]. The largest asymmetries are obtained at the lowest values of $\sqrt{S}$, where all the asymmetries are dominated by the gluon asymmetry.

For pp collisions, neglecting the $\bar{q}q$ channel for reasons explained earlier, we find that

$$A_{pp} = \left[1 + \mathcal{O}(v^4)\right] \frac{\Delta\mathcal{L}_{gg}}{\mathcal{L}_{gg}} \quad ({}^1S_0, {}^1P_1 \text{ quarkonia}), \quad (3.4)$$

where the order v^4 corrections come from the $\bar{q}q$ channel for the 1S_0 resonance, and in addition from the order v^9 matrix elements for the 1P_1 resonance.

The case of 3P_J states is very interesting phenomenologically. For $\gamma p \rightarrow \chi_J$ reactions, we need to retain only the contribution from the colour amplitude D . The cross sections then begin at order v^9 , and we find that

$$A_{\gamma p} = \frac{\Delta g(x)}{g(x)} \frac{\tilde{\Theta}_D^{\chi_J}(9)}{\Theta_D^{\chi_J}(9)} \quad ({}^3P_J \text{ quarkonia}). \quad (3.5)$$

Since both the non-perturbative matrix elements involved are unknown, it might seem that there is no predictive power to the model. However, using heavy quark symmetry, and the scaling form of eq. (3.1) it appears that $\tilde{\Theta}_D^{\chi_J}(9) = 0$. As a result, we obtain the prediction

$$A_{\gamma p}^{\chi_0} = A_{\gamma p}^{\chi_1} = A_{\gamma p}^{\chi_2} \approx [0 + \mathcal{O}(v^2)] \frac{\Delta g(x)}{g(x)}, \quad (3.6)$$

where the order v^2 terms come from the un-calculated order v^{11} terms in the asymmetry and from corrections to heavy quark spin symmetry.

We can make similar arguments for the polarisation asymmetries in $\gamma\gamma$ collisions. We find that

$$A_{\gamma\gamma}^{\chi_0} = 1 + \mathcal{O}(v^4) = -A_{\gamma\gamma}^{\chi_2}. \quad (3.7)$$

The case of χ_1 is more complicated, since a ratio of two unknown matrix elements is involved. However, using heavy-quark spin symmetry this reduces to

$$A_{\gamma\gamma}^{\chi_1} = -\frac{1}{7} + \mathcal{O}(v^2). \quad (3.8)$$

Unfortunately, no polarised $\gamma\gamma$ experiments have been planned to test this simple prediction of NRQCD and heavy quark spin symmetry.

A straightforward application of the NRQCD scaling laws would lead us to the conclusion that the asymmetries for $pp \rightarrow \chi_{0,2}$ are given by

$$A_{pp}^{\chi_0} \approx -A_{pp}^{\chi_2} \approx \frac{\Delta\mathcal{L}_{gg}}{\mathcal{L}_{gg}} + \mathcal{O}(v^4), \quad (3.9)$$

where we have neglected the contribution of the $\bar{q}q$ channel. The asymmetry for χ_1 production is

$$A_{pp}^{\chi_1} = \frac{\tilde{\Theta}_S^{\chi_1}(9) + \tilde{\Theta}_D^{\chi_1}(9) + \tilde{\Theta}_F^{\chi_1}(9)}{\Theta_S^{\chi_1}(9) + \Theta_D^{\chi_1}(9) + \Theta_F^{\chi_1}(9)} \left[\frac{\Delta\mathcal{L}_{gg}}{\mathcal{L}_{gg}} \right]. \quad (3.10)$$

The ratio of matrix elements can be estimated using heavy quark spin symmetry and the scaling relations in eq. (3.1). $\tilde{\Theta}_D^{\chi_J}(9)$ vanishes in this approximation and the terms $\tilde{\Theta}_{S,F}^{\chi_J}(9)$ come with opposite signs. The numerator is positive but small and we expect—

$$A_{pp}^{\chi_1} \approx 0.2 \frac{\Delta\mathcal{L}_{gg}}{\mathcal{L}_{gg}}. \quad (3.11)$$

The $\bar{q}q$ channel remains negligible even at $\sqrt{S} = 500$ GeV.

The ψ' asymmetry is straightforward. If the $\bar{q}q$ channel is neglected then

$$A_{pp}^{\psi'} = \frac{\Delta\mathcal{L}_{gg}}{\mathcal{L}_{gg}} \frac{\tilde{\Theta}_D^{\psi'}(7)}{\Theta_D^{\psi'}(7)} + \mathcal{O}(v^2), \quad A_{\gamma p}^{\psi'} = \frac{\Delta g(x)}{g(x)} \frac{\tilde{\Theta}_D^{\psi'}(7)}{\Theta_D^{\psi'}(7)} + \mathcal{O}(v^2), \quad (3.12)$$

and the form of the non-perturbative matrix elements is the same as that for J/ψ . The numerator can be shown to vanish using heavy quark spin symmetry and eq. (3.1). A numerical estimate of the subleading order v^9 terms shows that the ratio of non-perturbative matrix elements is small (of the order of 0.05). At low $\sqrt{S}$ the asymmetry is dominated by the asymmetry in the gluon densities. However, at $\sqrt{S}$ of a few hundred GeV, the gluon asymmetry reduces to about 0.001, and the neglected $\bar{q}q$ channel becomes important. These terms appear with the opposite sign and reduce the asymmetry even further. As a consequence, we expect the ψ' asymmetry to be small for all $\sqrt{S}$ —

$$A_{pp}^{\psi'} \approx 0, \quad A_{\gamma p}^{\psi'} \approx 0. \quad (3.13)$$

Any attempt to see structure in this asymmetry would probably require unreasonably large statistics.

The J/ψ asymmetry seems to be enormously complicated because of the feed-down from radiative decays of the χ states. However, a major simplification occurs because of the near vanishing asymmetry in direct J/ψ production (the argument is the same as for ψ'). Thus the asymmetry comes entirely from the 20–40% of the cross section due to χ decays. Taking into account the ratios of the production cross sections of χ and the branching fractions for their decays into J/ψ , we find that the χ_1 and χ_2 states contribute equally to J/ψ . Hence the J/ψ polarisation asymmetry is expected to be approximately

$$A_{pp}^{J/\psi} \approx -(0.15 \pm 0.05) \frac{\Delta \mathcal{L}_{gg}}{\mathcal{L}_{gg}}. \quad (3.14)$$

We summarize the predictions made on the basis of the NRQCD scaling in eq. (3.1) and the assumption of R_H depending only on the hadron H —

$$\begin{aligned} A_{pp}^{\psi'} &\ll -A_{pp}^{J/\psi} \approx A_{pp}^{\chi_1} < A_{pp}^{\chi_0} = -A_{pp}^{\chi_2} = \frac{\Delta \mathcal{L}_{gg}}{\mathcal{L}_{gg}}, \\ A_{\gamma p}^{\psi'} &\approx A_{\gamma p}^{J/\psi} \approx A_{\gamma p}^{\chi_1} \approx A_{\gamma p}^{\chi_0} \approx A_{\gamma p}^{\chi_2} \approx 0. \end{aligned} \quad (3.15)$$

The asymmetry for J/ψ in almost-elastic photo-production vanishes because the direct as well as feed-down contributions vanish.

Alternative points of view can be taken about the scaling of the non-perturbative matrix elements. These proceed on the assumption that the

mass scale, Λ , appearing in the soft matrix elements are smaller than m . If the scale Λ in eq. (3.1) is taken to be the QCD scale Λ_{QCD} [7], then the relation $v \sim \alpha_s(mv)$ [2] implies that $z \equiv \Lambda_{QCD}^2/m^2 < v^4$ for all m . The relative importance of terms then changes dramatically, since the lowest dimensional operators dominate. Since the dimension 4 terms $\mathcal{O}_8^{\chi_J}({}^1S_0^0)$ appear in the χ cross sections at order $v^9 > zv^5$, they dominate the cross section. Heavy quark spin symmetry then gives the χ_1/χ_2 ratio to be $5/3$, in agreement with the pion-nucleon observations. The same χ_1/χ_2 ratio is then predicted for charm and bottom mesons in hadro- and photo-production, and the result remains stable even when higher order terms are introduced. The rearrangement of the series also leads to $\Theta_D^{J/\psi}(7) \approx \tilde{\Theta}_D^{J/\psi}(7) \neq 0$ and $\Theta_D^{\chi_J}(9) \approx \tilde{\Theta}_D^{\chi_J}(9) \neq 0$. The consequences for the asymmetries are—

$$\begin{aligned} A_{pp}^{J/\psi} &\approx A_{pp}^{\psi'} \approx A_{pp}^{\chi_0} \approx A_{pp}^{\chi_1} \approx A_{pp}^{\chi_2} = \frac{\Delta\mathcal{L}_{gg}}{\mathcal{L}_{gg}}, \\ A_{\gamma p}^{J/\psi} &\approx A_{\gamma p}^{\psi'} \approx A_{\gamma p}^{\chi_0} \approx A_{\gamma p}^{\chi_1} \approx A_{\gamma p}^{\chi_2} = \frac{\Delta g(x)}{g(x)}, \end{aligned} \quad (3.16)$$

and totally different from the NRQCD results summarized in eqs. (3.15). These differences make low-energy double polarised experiments a good testing ground for NRQCD, and the entire phenomenology of low-energy quarkonium production. Other choices of the mass scale Λ have also been suggested [18]. These give somewhat different results. The asymmetries for $\eta_{c,b}$ and $h_{c,b}$ are independent of such assumptions, and hence form good measures of polarised gluon densities.

In conclusion, low energy double polarised asymmetries are a good test-bed for understanding the origin of all observed systematics in fixed target hadro- and photo-production of charmonium. The high order computations presented here provide us with a set of processes which can be used to test aspects of NRQCD factorisation and scaling. In addition, 1S_0 and 1P_1 charmonium and bottomonium states are well suited for measurements of polarised gluon densities. Assuming standard NRQCD power counting laws, higher order terms are likely to be phenomenologically important for double polarisation asymmetries of 3P_1 and excited 3S_0 states in both hadro- and photo-production.

References

- [1] W.E. Caswell and G.P. Lepage, *Phys. Lett.*, B 167 (1986) 437.
- [2] G. T. Bodwin, E. Braaten and G. P. Lepage, *Phys. Rev.*, D 51 (1995) 1125; erratum *ibid.*, D 55 (1997) 5853.
- [3] E. Braaten, M. A. Doncheski, S. Fleming and M. Mangano, *Phys. Lett.*, B 333 (1994) 548;
D. P. Roy and K. Sridhar, *Phys. Lett.*, B 339 (1994) 141;
M. Cacciari and M. Greco, *Phys. Rev. Lett.*, 73 (1994) 1586.
- [4] S. Gupta and K. Sridhar, *Phys. Rev.*, D 54 (1996) 5545.
- [5] M. Beneke and I. Rothstein, *Phys. Rev.*, D 54 (1996) 2005;
M. Beneke, preprint CERN-TH/97-55, hep-ph/9703429.
- [6] S. Gupta and K. Sridhar, *Phys. Rev.*, D 55 (1997) 2650.
- [7] S. Gupta and P. Mathews, preprint hep-ph/9703370, to appear in *Phys. Rev.*, D.
- [8] S. Gupta and P. Mathews, *Phys. Rev.*, D 55 (1997) 7144.
- [9] O. Terayev and A. Tkabladze, preprint hep-ph/9612301.
- [10] E. Braaten and Y. Chen, *Phys. Rev.*, D 54 (1996) 3216.
- [11] M. Glück, E. Reya and M. Stratmann and W. Vogelsang, *Phys. Rev.*, D 53 (1996) 4775.
- [12] Yu. L. Dokshitzer, D. I. Dyakonov and S. I. Troyan, *Phys. Rep.*, 58 (1980) 269.
- [13] M. Beneke and M. Kramer, *Phys. Rev.*, D 55 (1997) 5269.
- [14] K. Hagan-Ingram (E771 Collaboration), unpublished; data quoted in the talk by M. Beneke [5].
- [15] D. A. Bauer *et al.*, *Phys. Rev. Lett.*, 54 (1985) 753.
- [16] V. Koreshev *et al.*, *Phys. Rev. Lett.*, 77 (1996) 4294.

- [17] M. Klein, preprint DESY 96-218, hep-ph/9610398;
B. Clerbaux, preprint DESY-96-162F;
S. Schiek *et al.*, H1 Collaboration, preprint DESY-96-162E;
S. Aid *et al.*, H1 Collaboration, *Nucl. Phys.*, B 472 (1996) 3.
- [18] G. Schuler, preprint CERN-TH-97-012, hep-ph/9702230.